

兔抗 HDAC5 多克隆抗体

中文名称：兔抗 HDAC5 多克隆抗体

英文名称：Anti-HDAC5 rabbit polyclonal antibody

别名：HD5; NY-CO-9

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：HDAC5

反应种属：Human, Mouse

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been found for this gene.
--------------------	---

Applications:	ELISA, IHC
Name of antibody:	HDAC5
Immunogen:	Synthetic peptide of human HDAC5
Full name:	histone deacetylase 5
Synonyms :	HD5; NY-CO-9
SwissProt:	Q9UQL6
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human gastric cancer and Human tonsil
IHC Recommend dilution :	25-100

