

兔抗 ERBB3 多克隆抗体

- 中文名称: 兔抗 ERBB3 多克隆抗体
- 英文名称: Anti-ERBB3 rabbit polyclonal antibody

别 名: HER3, LCCS2, ErbB-3, c-erbB3, erbB3-S, MDA-BF-1, c-erbB-3, p180-ErbB3, p45-sErbB3, p85-sErbB3

- 抗 原: ERBB3
- 储存: 冷冻 (-20℃)
- 宿 主: Rabbit
- 相关类别: 一抗
- 反应种属: Human, Mouse, Rat
- 标记物: Unconjugate
- 克隆类型: rabbit polyclonal

技术规格

Background:	This gene encodes a member of the epidermal growth f actor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore ca n bind this ligand but not convey the signal into the ce II through protein phosphorylation. However, it does for m heterodimers with other EGF receptor family member s which do have kinase activity. Heterodimerization lead s to the activation of pathways which lead to cell prolif eration or differentiation. Amplification of this gene and/ or overexpression of its protein have been reported in n
	or overexpression of its protein have been reported in n umerous cancers, including prostate, bladder, and breast



	tumors. Alternate transcriptional splice variants encoding different isoforms have been characterized. One isoform lacks the intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have a lso been reported, but they have not been thoroughly c haracterized.
Applications:	ELISA, IHC
Name of antibody:	ERBB3
Immunogen:	Fusion protein of human ERBB3
Full name:	erb-b2 receptor tyrosine kinase 3
Synonyms :	HER3, LCCS2, ErbB-3, c-erbB3, erbB3-S, MDA-BF-1, c-erb B-3, p180-ErbB3, p45-sErbB3, p85-sErbB3
SwissProt:	P21860
ELISA Recommended dilution:	1000-5000
IHC positive control:	Human colon cancer
IHC Recommend dilution:	50-150

