

兔抗 RCAN2 多克隆抗体

中文名称：兔抗 RCAN2 多克隆抗体

英文名称：Anti-RCAN2 rabbit polyclonal antibody

别名：CSP2; RCN2; MCIP2; ZAKI4; ZAKI-4; DSCR1L1

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：RCAN2

反应种属：Human, Mouse, Rat

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|----------------------|---|
| Background: | This gene encodes a member of the regulator of calcineurin (RCAN) protein family. These proteins play a role in many physiological processes by binding to the catalytic domain of calcineurin A, inhibiting calcineurin-mediated nuclear translocation of the transcription factor NFATC1. Expression of this gene in skin fibroblasts is upregulated by thyroid hormone, and the encoded protein may also play a role in endothelial cell function and angiogenesis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. |
| Applications: | ELISA, IHC |

| | |
|------------------------------------|---|
| Name of antibody: | RCAN2 |
| Immunogen: | Synthetic peptide of human RCAN2 |
| Full name: | regulator of calcineurin 2 |
| Synonyms : | CSP2; RCN2; MCIP2; ZAKI4; ZAKI-4; DSCR1L1 |
| SwissProt: | Q14206 |
| ELISA Recommended dilution: | 2000-5000 |
| IHC positive control: | Human brain |
| IHC Recommend dilution: | 25-100 |

