

兔抗 FLT3(Phospho-Tyr842) 多克隆抗体

- 中文名称: 兔抗 FLT3(Phospho-Tyr842) 多克隆抗体
- 英文名称: Anti-FLT3(Phospho-Tyr842) rabbit polyclonal antibody
- 别名: FLK2; STK1; CD135; FLK-2
- 相关类别: 一抗
- 储存: 冷冻(-20℃)避光
- 宿 主: Rabbit
- 抗原: FLT3(Phospho-Tyr842)
- 反应种属: Human Mouse
- 标记物: Unconjugate
- 克隆类型: Unconjugate

技术规格

| Background: | This gene encodes a class III receptor tyrosine kinas e that regulates hematopoiesis. The receptor consist s of an extracellular domain composed of five immu noglobulin-like domains, one transmembrane region, and a cytoplasmic kinase domain split into two part s by a kinase-insert domain. The receptor is activate d by binding of the fms-related tyrosine kinase 3 li gand to the extracellular domain, which induces ho modimer formation in the plasma membrane leadin g to autophosphorylation of the receptor. |
|-------------------|---|
| Applications: | WB |
| Name of antibody: | FLT3(Phospho-Tyr842) |
| Immunogen: | Peptide sequence around phosphorylation site of tyr |



全国订货电话 4008-723-722

| | osine 842(S-N-Y(p)-V-V) derived from Human FLT3 . |
|--------------------------|---|
| Full name: | fms-related tyrosine kinase 3 |
| Synonyms : | FLK2; STK1; CD135; FLK-2 |
| SwissProt: | P36888 |
| WB Predicted band size: | 113 kDa |
| WB Positive control: | HepG2 cells lysates treated with EGF |
| WB Recommended dilution: | 500-1000 |

